



Give Your Patients a **Better Way** to Manage Their Pain

NON-PHARMACOLOGICAL ALTERNATIVES FOR:

Postoperative Pain and Swelling
Chronic and Acute Pain

REDUCE OPIOID USAGE

Non-pharmacological treatment options give your patients a better opportunity to use less opioids to manage pain.

ZYNEX REPRESENTATIVE:

Rep Name

Phone: (800) 495-6670

Fax: +1 (800) 495-6695

Email: repname@zynex.com



SPECIALIZING IN TRICARE AND WORKER'S COMP



1



2



3

EZ PRESCRIBE™

STEP 1
Complete Rx

STEP 2
Give Handout to Patient

STEP 3
Fax Rx & Demographics to +1 (800) 495-6695

ZYNEX WILL TAKE CARE OF THE REST



PRESCRIPTION & LETTER OF MEDICAL NECESSITY

PRESCRIBER'S PRINTED NAME* NPI*

CLINIC ADDRESS CITY ST. ZIP

PHONE FAX

IF THIS DEVICE WAS REQUESTED BY ANOTHER MEDICAL PROVIDER, PLEASE INCLUDE THEIR NAME/ADDRESS/ZIP HERE

PATIENT NAME* DOB* DATE OF INJURY DATE OF SURGERY

PRIMARY PHONE* SECONDARY PHONE/ EMAIL PRIMARY LANGUAGE (IF NOT ENGLISH)

INSURANCE TYPE
 Work Comp TRICARE Auto Attorney/PI Commercial Medicare/Medicaid Uninsured
(e.g.- UHC,BCBS,Cigna,Aetna,etc.)

ATTORNEY/ADJUSTER PHONE/EMAIL CASE/CLAIM NUMBER

E-STIM

Zynex InWave + Probe
Length of Need: Lifetime of Treatment 3-18 Months (if unchecked = Lifetime of Treatment)
DX Codes: N39.41 Urge IC N39.3 Stress IC
 N39.46 Mixed IC Other _____

Substitution for this device is NOT ALLOWED without my written approval

BRACING (FILL OUT PATIENT EQUIPMENT PROTOCOL ON BACK OF PATIENT HANDOUT)

Zynex Pro Hybrid Lumbosacral Orthosis Zynex Pro Wrist DX Code(s) _____
 Zynex Pro Lumbosacral Orthosis Zynex Pro Post-Op Knee Knee: Left Right
 Zynex Pro Low-Profile Lumbosacral Orthosis Zynex Pro OA Knee with Suspension Wrap
 Zynex Pro Thoracic-Lumbosacral Orthosis

Please document in patient's chart the reason(s) for prescribing and fax with Rx

CERVICAL TRACTION (FILL OUT PATIENT EQUIPMENT PROTOCOL ON BACK OF PATIENT HANDOUT)

Cervical Traction (Saunders or ComforTrac) DX Code(s) _____

HOT/COLD THERAPY (FILL OUT PATIENT EQUIPMENT PROTOCOL ON BACK OF PATIENT HANDOUT)

Zynex CryoHeat circulating pump & therapy blanket Length of Need: 15 Days 30 Days 45 Days Other ___ Days
 Zynex DynaComp cold compression & therapy blanket DX Code(s) _____

I have reviewed the hot/cold therapy contraindications on the back of this prescription prior to prescribing for patient.

PRESCRIBER'S SIGNATURE* DATE*

I CERTIFY THAT THE EQUIPMENT AND SUPPLIES I PRESCRIBED ARE MEDICALLY NECESSARY FOR THIS PATIENT'S WELL-BEING; THIS IS NOT PRESCRIBED AS CONVENIENCE EQUIPMENT. IN MY PROFESSIONAL OPINION, THE EQUIPMENT IS BOTH REASONABLE AND NECESSARY IN REFERENCE TO THE ACCEPTED STANDARDS OF MEDICAL PRACTICE AND TREATMENT FOR THIS PATIENT'S CONDITION. SUBSTITUTION FOR THIS DEVICE IS NOT ALLOWED WITHOUT MY WRITTEN APPROVAL.

Additional Device Information

Cervical Traction

Our Home Cervical Traction Devices are medically necessary because the patient requires greater than 20 pounds of force.

Please Review Hot/Cold therapy contraindications prior to prescribing for patient

CONTRAINDICATION - WHEN NOT TO USE HOT THERAPY

DO NOT use hot therapy on patients with any of the following contraindications:

- History of heat injury or adverse reactions to hot therapy
- Incoherence due to sedation, anesthesia, coma or sleep
- Decreased skin sensitivity or localized circulation or wound healing problems
- Arteriosclerosis or atherosclerosis
- Localized malignancy in treatment area
- Hypersensitivity to heat
- Local tissue infection, swelling or inflammation

DO NOT APPLY ABDOMINALLY IF PREGNANT

CONTRAINDICATION - WHEN NOT TO USE COLD THERAPY

DO NOT use cold therapy on patients with any of the following contraindications:

- History of cold injury, frostbite or adverse reactions to cold therapy
- Incoherence due to sedation, anesthesia, coma or sleep
- Decreased skin sensitivity or localized circulation or wound healing problems, including those caused by multiple surgical procedures
- Circulatory syndromes due to Raynaud's disease, Buerger's disease, peripheral vascular disease, vasospastic disorders, or hypercoagulable clotting disorders
- Hand/wrist or foot/ankle surgery exhibiting polyneuropathy
- Hypersensitivity to cold
- Local tissue infection
- Diabetes

DO NOT USE ON CHILDREN UNDER 12 YEARS OF AGE

Please Fax To: +1 (800) 495-6695

Include Patient Demographics

****Please include patient chart notes supporting Rx****

IMPORTANT **GIVE TO PATIENT**

HERE'S HOW TO OBTAIN YOUR PRESCRIBED MEDICAL DEVICE

Step 1

- Zynex will text and call you
(usually the same day we receive your Rx!)



Step 2

- Have your medical insurance



Step 3

- Zynex will ship your device



You can learn how to use the NexWave by watching the instructional video at
<http://www.zynex.com/nw>

*If you have not heard from us within 24 hours, please call Patient Support
at (800) 495-6670. You may also go online to complete your order at
www.zynex.com/ship or by scanning here:*



(SEE BACK FOR: EQUIPMENT USE)



Equipment Use Protocol

Spinal Bracing Equipment Use Protocol

Pain Management Use

Wear ___ min./day and increase wear time by ___ min. every ___ day(s)

Wear spinal brace while performing the following:

Sitting Standing Walking Exercising Other _____

Instructions: _____

Post Operative Use

Day ___ through Day ___ Wear continuously with a break of ___ min. every ___ hour(s)

Day ___ through Day ___ Wear continuously with a break of ___ min. every ___ hour(s)

Wear spinal brace while performing the following:

Sitting Standing Walking Exercising Other _____

Instructions: _____

Knee Bracing Equipment Use Protocol

Range of Motion Restriction: _____

Offloading: Medial Lateral

Instructions: _____

Cervical Traction Equipment Use Protocol (Tx = Treatment)

Choose Degree of Flexion: 10° 15° 20°

Tension Type: Sustained ___ lbs. of Force ___ min./Tx ___ Tx/Day

Tension Type: Intermittent ___ lbs. of Force ___ min. of ON Time ___ min. of OFF Time ___ Tx/day

Hot/Cold Therapy Equipment Use Protocol (Tx = Treatment)

Cold Therapy (40° - 50°):

Day ___ through Day ___ Use continuously with a break of ___ min. every ___ hour(s)

Day ___ through Day ___ Use: ___ min./Tx every ___ hour(s)

Use after exercise for ___ min./Tx

Hot Therapy: (80° - 120°)

Use after day number ___ Before exercise: ___ min./Tx

Loosen stiff muscles/Joint: ___ min./Tx

See Back For: "How to Obtain Your Prescribed Device"